

# BIOMIN<sup>®</sup> BORON

## Mineral for Soil and Foliar Application

**1-0-0**

### Product Information

**BIOMIN BORON** contains boron for soil and/or foliar application. Completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. Designed to prevent and correct boron deficiency and to boost crops during critical or fast growing periods. May be applied with nitrogen fertilizers.

#### GUARANTEED ANALYSIS

Total Nitrogen (N).....	1.0%
1.0% Water Soluble Nitrogen	
Boron (B).....	3.0%

Derived from sodium tetraborate and soy protein hydrolysate.

**KEEP OUT OF REACH OF CHILDREN**

**SHAKE WELL BEFORE USE  
PREVENTS AND CORRECTS  
MINERAL DEFICIENCY**

**NET CONTENTS:** 2.5 gal. (9.46 l)  
Weight per Gallon: 9.5 lb. (4.3 kg) @ 68°F  
**U.S. Pat.# 5,504,055**  
**Lot #:**



**MANUFACTURED BY: JH BIOTECH, INC.**  
P.O. Box 3538, Ventura, CA 93006 U.S.A.  
805-650-8933 [jhbiotech.com](http://jhbiotech.com)

### Directions and General Recommendations

#### SOIL APPLICATION

**BIOMIN BORON** can be applied before or at planting through irrigation water, as banding or broadcast. The application rate is 1 qt. to 1 gal./ac. (2.5-10 l/ha) depending on crop, soil conditions and application methods. For maintenance, 1 qt./ac. (2.5 l/ha) is recommended. Dilute **BIOMIN BORON** with enough water to cover entire acre.

#### FOLIAR APPLICATION

**FIELD AND ROW CROPS:** Apply 1-4 pt./ac. (1.25-5 l/ha) when crops are 6-12 in. (15-30 cm) in high or when deficiency symptoms appear. Repeat application 2-4 weeks later.

**FRUIT TREES AND GRAPES:** Apply 1-4 pt./ac. (1.25-5 l/ha) starting at bud break or when deficiency symptoms appear. Repeat the application 2-4 weeks later.

**VEGETABLES AND POTATOES:** Apply 1-4 pt./ac. (1.25-5 l/ha) at about 2-4 weeks after emergence or if deficiency symptoms appear. Repeat the application 2-3 weeks later. In greenhouses and plastic tunnels, application may be repeated every week or every other week.

**ALMONDS, WALNUTS, PISTACHIOS, AND OTHER NUT CROPS:** Apply 1-4 pt./ac. (1.25-5 l/ha) before bloom. Repeat the application if necessary.

For best results, apply when plants are in an active growing stage, after irrigation or rainfall. For foliar application, dilute it with at least 100 times of water.

**BIOMIN BORON** can be applied to fruit trees, field crops, vegetables, greenhouse crops, potatoes, grapes, citrus, and ornamental gardens.

**CAUTION:** This product contains Boron. Do not apply to Boron sensitive crops. Use of the product on any crop other than those recommended may result in serious injury to the crop(s).

**Precautionary statement:** do not swallow. Wash hands thoroughly after handling. **First Aid:** if swallowed; seek medical attention if feeling unwell. Rinse mouth. **Storage & Disposal:** store in a safe manner. Dispose product/container in accordance with federal, state, and local regulations.

#### LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling, or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

051001-D-B-2

Copyright © 2019 All rights reserved