# FLORAPRO® POWDER FEED PROGRAM | HIGH EC

| FLORAPRO® POWDER HIGH EC STOCK SOLUTION FEED PROGRAM Feed rates in mL/gal | | Propagation | Vegetative | Early Flower | Mid Flower | Late Flower | Finish |
|---|---|---|---|---|---|---|---|
| Use 1.0 lb per gallon concentrated stock solutions | ← Mixing Order | | | | | | |
| | FloraPro® Ca + Micros | 7.0 | 31.0 | 30.0 | 28.5 | 19.0 | 12.5 |
| | FloraPro® Grow | 9.0 | 38.0 | – | – | – | – |
| | FloraPro® Bloom | – | – | 33.0 | 25.5 | 15.0 | – |
| | FloraPro® Late Bloom | – | – | 12.5 | 21.0 | 26.0 | 32.0 |
| | TrueFlo (mL/gal) | 3 | 4 | 4 | 4 | 4 | 4 - 9 |
| | Nitrogen (ppm) | 60 | 250 | 210 | 185 | 115 | 50 |
| | Target pH | 5.5 - 5.8 | 5.6 - 6.1 | 5.7 - 6.2 | | | |
| | EC (mS/cm) | 0.6 - 0.8 | 2.5 - 2.7 | 2.6 - 2.8 | 2.4 - 2.6 | 1.9 - 2.1 | 1.5 - 1.7 |

Mixing chamber is strongly recommended between the injection of Ca+Micros and the other base inputs

Coco & Rockwool

Learn how to prepare and verify a stock solution
(Video Guide)

| FLORAPRO® POWDER HIGH EC STOCK SOLUTION FEED PROGRAM Feed rates in mL/gal | | Propagation | Vegetative | | | Early Flower | | | Mid Flower | | | Late Flower | | Finish |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Wk 1 | Wk 1 | Wk 2 | Wk 3 | Wk 1 | Wk 2 | *Wk 3 | Wk 4 | Wk 5 | Wk 6 | Wk 7 | Wk 8 | Wk 9 |
| Use 1.0 lb per gallon concentrated stock solutions | FloraPro® Ca + Micros | 7.0 | 31.0 | 31.0 | 31.0 | 30.0 | 30.0 | 30.0 | 28.5 | 28.5 | 28.5 | 19.0 | 19.0 | 12.5 |
| | FloraPro® Grow | 9.0 | 38.0 | 38.0 | 38.0 | – | – | – | – | – | – | – | – | – |
| | FloraPro® Bloom | – | – | – | – | 33.0 | 33.0 | 33.0 | 25.5 | 25.5 | 25.5 | 15.0 | 15.0 | – |
| | FloraPro® Late Bloom | – | – | – | – | 12.5 | 12.5 | 12.5 | 21.0 | 21.0 | 21.0 | 26.0 | 26.0 | 32.0 |
| | TrueFlo (mL/gal) | 3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 - 9 |
| | Nitrogen (ppm) | 60 | 250 | 250 | 250 | 210 | 210 | 210 | 185 | 185 | 185 | 115 | 115 | 50 |
| | Target pH | 5.5 - 5.8 | 5.6 - 6.1 | | | 5.7 - 6.2 | | | | | | | | |
| | EC (mS/cm) | 0.6 - 0.8 | 2.5 - 2.7 | | | 2.6 - 2.8 | | | 2.4 - 2.6 | | | 1.9 - 2.1 | | 1.5 - 1.7 |

*For an 8-week flowering schedule, reduce the 'Early Flower' phase by one week.



Disclaimer: Feed chart provided as recommendation only. Actual plant needs and/or results achieved may vary dependent upon your specific cultivation conditions and protocols.

©2026 General Hydroponics.
GH_FloraProFeedchartUSC-HighEC_261029ae

# FLORAPRO® POWDER FEED PROGRAM | HIGH EC w/FUL+

| FLORAPRO® POWDER HIGH EC w/FUL+ STOCK SOLUTION FEED PROGRAM<br>Feed rates in mL/gal | | Propagation | Vegetative | Early Flower | Mid Flower | Late Flower | Finish |
|---|---|---|---|---|---|---|---|
| Use 1.0 lb per gallon concentrated stock solutions | ← Mixing Order | | | | | | |
| | FloraPro® Ca + Micros | 7.0 | 31.0 | 30.0 | 28.5 | 19.0 | 12.5 |
| | FloraPro® Grow | 9.0 | 38.0 | – | – | – | – |
| | FloraPro® Bloom | – | – | 33.0 | 25.5 | 15.0 | – |
| | FloraPro® Late Bloom | – | – | 12.5 | 21.0 | 26.0 | 32.0 |
| | FloraPro® FUL+ | – | 2 | 2 | 4 | 4 | 4 |
| | TrueFlo | 3 | 5 | 5 | 5 | 5 | 5 - 9 |
| | Nitrogen (ppm) | 60 | 255 | 215 | 195 | 125 | 60 |
| | Target pH | 5.5 - 5.8 | 5.6 - 6.1 | 5.7 - 6.2 | | | |
| | EC (mS/cm) | 0.6 - 0.8 | 2.6 - 2.8 | 2.7 - 2.9 | 2.6 - 2.8 | 2.1 - 2.3 | 1.7 - 1.9 |

Mixing chamber is strongly recommended between the injection of Ca+Micros and the other base inputs

Coco & Rockwool

Learn how to prepare and verify a stock solution
(Video Guide)

| FLORAPRO® POWDER HIGH EC w/FUL+ STOCK SOLUTION FEED PROGRAM<br>Feed rates in mL/gal | | Propagation | Vegetative | | | Early Flower | | | Mid Flower | | | Late Flower | | Finish |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Wk 1 | Wk 1 | Wk 2 | Wk 3 | Wk 1 | Wk 2 | Wk 3 | Wk 4 | Wk 5 | Wk 6 | Wk 7 | Wk 8 | Wk 9 |
| Use 1.0 lb per gallon concentrated stock solutions | FloraPro® Ca + Micros | 7.0 | 31.0 | 31.0 | 31.0 | 30.0 | 30.0 | 30.0 | 28.5 | 28.5 | 28.5 | 19.0 | 19.0 | 12.5 |
| | FloraPro® Grow | 9.0 | 38.0 | 38.0 | 38.0 | – | – | – | – | – | – | – | – | – |
| | FloraPro® Bloom | – | – | – | – | 33.0 | 33.0 | 33.0 | 25.5 | 25.5 | 25.5 | 15.0 | 15.0 | – |
| | FloraPro® Late Bloom | – | – | – | – | 12.5 | 12.5 | 12.5 | 21.0 | 21.0 | 21.0 | 26.0 | 26.0 | 32.0 |
| | FloraPro® FUL+ | – | 2 | 2 | 2 | 2 | 2 | 2 | 4 | 4 | 4 | 4 | 4 | 4 |
| | TrueFlo | 3 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 - 9 |
| | Nitrogen (ppm) | 60 | 255 | 255 | 255 | 215 | 215 | 215 | 195 | 195 | 195 | 125 | 125 | 60 |
| | Target pH | 5.5 - 5.8 | 5.6 - 6.1 | | | 5.7 - 6.2 | | | | | | | | |
| | EC (mS/cm) | 0.6 - 0.8 | 2.6 - 2.8 | | | 2.7 - 2.9 | | | 2.6 - 2.8 | | | 2.1 - 2.3 | | 1.7 - 1.9 |



Disclaimer: Feed chart provided as recommendation only. Actual plant needs and/or results achieved may vary dependent upon your specific cultivation conditions and protocols.

©2026 General Hydroponics.
GH_FloraProFeedchartUSC-Ful_260114ae